

# Comparative Phytochemical and Antibacterial Studies on the bark of *Alstonia scholaris* R.Br. and *Alstonia macrophylla* Wall. ex G.Don

Khyade M. S.<sup>1\*</sup> and Vaikos N. P.

<sup>1</sup>Post Graduate Department of Botany, Sangamner Nagarpalika Arts, D.J. Malpani Commerce and B.N. Sarda Science College, Sangamner, Dist. Ahmednagar-422 605 (MS), India.

<sup>2</sup>Department of Botany, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad-431004 (MS), India

## ABSTRACT

Comparative phytochemical and antibacterial activities of bark of *Alstonia scholaris* and *Alstonia macrophylla* were investigated. The successive different solvent extracts showed the presence of alkaloids, phenolics, saponins and tannins were found in both the species. The antibacterial activities of bark of *A. scholaris* and *A. macrophylla* in successive different solvent were tested against gram +ve and gram -ve organisms. The chloroform extracts of *A. macrophylla* showed broader spectrum of antibacterial activity when compared with *A. scholaris*. However, *Alstonia scholaris* is widely used medicinal plant.

**Keywords:** Comparative, bark, *Alstonia* species, phytochemical, antibacterial

**Editor:** Dr. Mueen Ahmed K.K., Phcog.Net

**Copyright:** © 2010 Phcog.net

**\*Author for Correspondence:** Email: mskhyade@rediffmail.com

## INTRODUCTION

Green plants represent a reservoir of effective chemotherapeutics and can provide valuable source of natural antimicrobials (1, 2). Plant species which have one or more of its organs containing substances that can be used for therapeutic purpose are called medicinal plants (3). Plants have been used as medicinal agents from the earliest day of mans existence (4, 5) and has made it necessary to study them in details in order to discriminate the kinds employed for different purposes (6). In particular, the antimicrobial activities of plant extracts have formed the basis of many applications, including raw and processed food formation, pharmaceuticals, alternative medicine and natural therapies (7).

Now a day, infectious diseases continue to be the major concern for health institutions, pharmaceutical companies and Governments all over the world, especially with the current increasing trends of multidrug resistance among emerging –reemerging bacterial pathogens to the available modern drug or antibiotics (8, 9). It is therefore very necessary that the search for newer antibiotic serves to be a continued process. Plants are the cheapest and safer alternative source of antimicrobials (10, 11). Therefore, with increasing acceptance of traditional medicine as an

alternative form of health care, the screening of plants for active compounds has become very important.

*Alstonia scholaris* R. Br. (Apocynaceae), popularly known as “Saptarni” or “Devil’s tree”, and *A. macrophylla*, are used in the traditional systems of medicine. However, the former one is widely recognized medicinal plant.

The bark of *A. scholaris* is bitter, tonic, astringent, expectorant, alterative, anthelmintic, emmenagogue and galactagogue. It has proved valuable in fever, chronic diarrhea and in advanced stages of dysentery (12–15). Whereas the bark of *Alstonia macrophylla* is used for the same purpose as that of *Alstonia scholaris* (16). Decoction of the stem bark is effective in stomach ache, skin diseases and urinary infections (17). Moreover, phytochemistry on these two species studied earlier (18–21). The genus *Alstonia* has been the subject of antimicrobial activities (22, 23). This work, however, is designed to evaluate the comparative account of antibacterial activity of related species.

## MATERIAL AND METHODS

### COLLECTION OF PLANT MATERIALS

The fresh barks of *A. scholaris* R. Br. were collected in the month of October 2005 from forest area of Aurangabad

district (M.S.) and that of *Alstonia macrophylla* from the Botanical Garden of Dr. Babasaheb Ambedkar Marathwada University, Aurangabad. The plant species were identified with the help of Flora of Marathwada (24) and voucher specimens have been deposited at the Botany department of the university. Plant samples were washed, shade dried at room temperature for 15 days.

### **PREPARATION OF EXTRACTS AND PHYTOCHEMICAL SCREENING**

The dried plant material was pulverized into fine powder using a grinder (mixer). About 50 gm of powdered material was extracted in Soxhlet extraction apparatus successively with 250 ml of each of the following solvents. Petroleum ether, chloroform, acetone and methanol (25). The extracts obtained with each solvent were filtered through Whatman filter paper No. 1 and the respective solvents were evaporated (at 40°C) with the help of heating mantle. The sticky greenish-brown substances were obtained and stored in refrigerator and were suspended in dimethyl sulphoxide (DMSO) for prior to use (26).

Some of the extracts of each solvent were used for the qualitative phytochemical screening for the identification of the various classes of active chemical constituents, using standard prescribed methods (27–30). The positive tests were noted as weak (+), moderate (++), strong (+++) and absent (–).

### **TEST CULTURE**

The test bacteria used for the screening antimicrobial activity were *Staphylococcus aureus*, *Bacillus subtilis*, *Micrococcus luteus* MTCC 106 *Escherichia coli*, *Salmonella typhi*, *Pseudomonas aeruginosa* MTCC 2488, *Klebsiella planticola* and *Bacillus megaterium*. The cultures were obtained from Microbial Type Culture Collection (MTCC), IMTEC, Chandigarh, India. Cultures were maintained as nutrient agar slants in screw-capped bottles and stored at 4°C. All cultures were checked for viability and purity by regular plating. Test cultures were prepared by transferring a loop full of bacteria from stock culture nutrient broth and incubated at 37±1°C for 24 hours.

### **SCREENING FOR ANTIBACTERIAL PROPERTIES**

The antibacterial activities of the successive bark extracts of petroleum ether, chloroform, acetone and methanol of both the plant species were tested by Agar well diffusion method (31). The culture plates were prepared by pouring 20 ml of sterile nutrient agar. 1ml inoculum suspension was spread uniformly over the agar medium using sterile glass rod to get uniform distribution of bacteria. A sterile

cork borer (6 mm) was used to make wells in each plate for extracts. These plates were labeled and 100µl of each plant extracts (at concentration of 100mg/ml) was added aseptically into the well. Then the plates were incubated for 24 h at 37°C during which the activity was evidenced by the presence of zone of inhibition surrounding the well. Ampicillin (40 µg/ml) was used as standard antibiotics. Each test was repeated three times and the antibacterial activity was expressed as the mean of diameter of the inhibition zones (mm) produced by the plant extracts when compared to the controls.

### **RESULTS AND DISCUSSION**

The results of the phytochemical screening of the bark of both species of *Alstonia* are shown in Table 1. It indicates the presence of alkaloids, phenolics, saponins and tannins. The presence of steroids and terpenoids are indicated only in *Alstonia scholaris* while cardiac glycosides and flavonoids are found alone in *A. macrophylla*. However there were complete absence of anthraquinones and leucoanthocyanins in both species.

The antibacterial activities of *A. scholaris* and *A. macrophylla* in petroleum ether, chloroform, acetone and methanol extracts against bacteria examined in the current study and their potency were qualitatively assessed by the presence or absence of inhibition zones and zone diameter (Table 2). The results showed that the extracts of *A. scholaris* mediated some degree of activity against bacteria. Except, *S. aureus* and *S. typhi* other strains like *M. luteus*, *B. subtilis*, *B. megaterium*, *K. planticola* *E. coli* and *P. aeruginosa* are inhibited by acetone extracts only, while other extracts such as petroleum ether, chloroform and methanol showed negative inhibition at test concentration.

The antibacterial activity of *A. macrophylla* extracts were the most prominent in activity against all bacteria tested at test concentration (Table 2). Except petroleum ether extracts, others such as chloroform, acetone and methanol showed broader spectrum of activity. However, chloroform extracts of *A. macrophylla* exhibited significant antibacterial activities against all the bacteria tested but, particularly, to *B. subtilis*, *S. aureus* and *E. coli* with a diameter same or greater than standard antibiotic ampicillin.

Based on these results, it is possible to conclude that of two species of *Alstonia*. *A. macrophylla* exhibited broader range of antibacterial activity to varying degrees, as it is a less known medicinal plant in the Indian literature. Particularly, the chloroform extracts of *A. macrophylla* showed significant antibacterial activities and could be used as antimicrobial agents in new drugs for therapy.

**Table -1: Phytochemical constituents of bark extracts of *A.scholaris* and *A. macrophylla***

Phytochemicals	<i>Alstonia scholaris</i>				<i>Alstonia macrophylla</i>			
	A	B	C	D	A	B	C	D
Alkaloids	—	+++	—	+++	—	+++	—	+++
Anthraquinones	—	—	—	—	—	—	—	—
Cardiac glycosides	—	—	—	—	—	—	++	+++
Coumarins	—	—	—	—	—	—	—	—
Flavonoids	—	—	—	—	—	++	++	+++
Leucoanthocyanins	—	—	—	—	—	—	—	—
Simple phenolics	—	—	+	++	—	++	++	+++
Steroids	—	—	—	++	—	—	—	—
Saponins	—	—	+	++	—	++	++	+++
Tannins	—	—	+	++	—	+	++	+++
Terpenoids	—	++	++	++	—	—	—	—

Note:- A - Pet ether, B - Chloroform, C - Acetone, D - Methanol, (+) - weak, (++) - moderate, (+++) - strong and (—) - absent.

**Table -2: Antibacterial activity of bark extracts of *A. scholaris* and *A. macrophylla***

Organisms	Gram stain + / -	Inhibition zone in diameters (mm / sensitive strains)									
		<i>A. scholaris</i>				<i>A. macrophylla</i>				DMSO	Ampicillin
		A	B	C	D	A	B	C	D		
<i>Staphylococcus aureus</i>	+	—	—	—	—	—	24	16	15	—	23
<i>Bacillus subtilis</i>	+	—	—	11	—	—	32	16	18	—	21
<i>Bacillus megaterium</i>	+	—	—	11	—	—	10	15	15	—	25
<i>Micrococcus luteus</i>	+	—	—	13	—	—	27	15	17	—	30
<i>Escherichia coli</i>	-	—	—	10	—	—	18	15	17	—	17
<i>Salmonella typhi</i>	-	—	—	—	—	—	11	13	10	—	19
<i>Pseudomonas aeruginosa</i>	-	—	—	10	—	—	—	14	14	—	16
<i>Klebsiella planticola</i>	-	—	—	11	—	—	22	18	15	—	21

Note:- A - Petroleum ether, B - Chloroform, C - Acetone, D - Methanol, — no inhibition, Figures are diameter of zone of inhibition.

## ACKNOWLEDGEMENT

The authors wish to thank Head of Botany Department, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad for providing the necessary laboratory facilities.

## REFERENCES

- Balandrin MF, Klocke JA, Wutule ES, Bollinger WH. Natural plant chemicals: Sources of industrial and medicinal materials. *Science* **228**:1154–1160 (1985).
- Satish H, Raveesha KA, Janardhana GR. Antibacterial activity of selected Peruvian medicinal plants. *J Ethnopharmacol* **88**:199–204 (1999).
- Sofowora A. *Medicinal plants and Traditional Medicine in Africa* (John Willy. New York, 1982) 289.
- Hansch C. *Comprehensive Medicinal Chemistry* Vol. **1**. (Pergamon Press, (1990)
- Shellard EJ. Medicines from plants. *Plant Med* **53**:6–8 (1987).
- Ghani A. *Medicinal Plants and Traditional Medicinal Portions: Problems and Prospects of their Standardization*. (The state of medicinal plant research in Nigeria, 1986) 65.
- Hammer KA, Carson CF, Riley TV. Antimicrobial activity of essential oils and other plant extracts. *App. Microbiol* **86**:985–990 (1999)
- Franklin TJ, Snow CA. *Biochemistry of antimicrobial action* 4<sup>th</sup> edn. (Chapman and Hall New York, 1989)134–155.
- Prescott L, Harley J, Klein DA. *Microbiology* 5th edn. (Mc-Graw Hill London, 2002) 820–950.
- Pretorius CJ, Watt E. Purification and identification of active components of *Carpobrotus edulis* L. *J Ethnopharmacol* **76**:87–91 (2001).
- Sharif MDM, Banik GR. Status and Utilization of Medicinal Plants in Rangamati of Bangladesh. *Res J Agri Biol Sci* **2**:268–273 (2006).
- Chopra RN, Nayar SI, Chopra IC. *Glossary of Indian Medicinal Plants*. (CSIR, New Delhi, India, 1956) 13–14.
- Dastur JF. *Medicinal plants of India and Pakistan* (Taraporewala and Co. Pvt. Ltd., Bombay, 1962)18.
- Joshi SG. *Medicinal plants* (Oxford and IBH publishing Co. Pvt. Ltd., New Delhi, 2000) 40–41.
- Siwakoti M, Siwakoti S. Ethnobotanical uses of plants among the star tribe of Nepal. In *Ethnobotany and medicinal plants of Indian subcon-*

- tinent*. (Maheshwari JK Editor, Scientific Publishers, Jodhpur, India, 2000) 99–108.
16. Anonymous. *The useful plants of India* (Publications and information Directorate, Council of Scientific and Industrial Research, New Delhi, 1992)
  17. Bhargava N. Ethanobotanical studies of the tribes of Andaman and Nicobar Islands, India. I onge. *Eco. Bot* **37**:110–119 (1983).
  18. Chakravarti D, Chakravarti RN, Ghose R. Triterpenes of *Alstonia scholaris* (dita-bark). *Experientia* **13**:277 (1957)
  19. Hadi Surya, Bremner John B. Initial studies on Alkaloids from Lombok Medicinal Plants. *Molecules* **6**:117–129 (2001).
  20. Kam TS, Nyeoh KT, Sim KM, Yoganathan K. Alkaloids from *Alstonia scholaris*. *Phytochemistry* **45**:1303–1305 (1997).
  21. Salim AA, Garson MJ, Craik DJ. New indole alkaloids from the bark of *Alstonia scholaris*. *J Nat Prod* **67**:1591–1594 (2004).
  22. Chattopadhyay D, Maiti K, Kundu AP, Chakraborty MS, Bharda R, Mandal SC, Mandal AB. Antimicrobial activity of *Alstonia macrophylla*: a folklore of bay islands. *J. Ethnopharmacology* **77**:49–55 (2001).
  23. Khan M, Omaloso R, Kihara M. Antibacterial activity of *Alstonia scholaris* and *Leea tetramera*. *Fitoterapia* **74**:736–40 (2003).
  24. Naik VN. *Flora of Marathwada* (Amrut Prakashan, Aurangabad (M.S), India, 1998) 528–529.
  25. Vogel AI. *Elementary practical organic chemistry* (2 edition), (Orient Longman Limited, 1988) 45–168.
  26. Beyer H, Walter W. *Organic chemistry*, 4<sup>th</sup> ed. (Published by Jerry March, 1997) 146–148.
  27. Harborne JB. *Phytochemical methods* (Chapman Hall London, 1984) 100–101.
  28. Trease E, Evans WC. 1987. *Pharmacognosy*, Billiare Tindall, London.
  29. Ajaiyeoba EO. Phytochemical and antimicrobial studies of *Gynandropsis gynandra* and *Buchholzia coriacea* extracts. *African Journal of Biomedical Research* **3**:161–165 (2000).
  30. Edeoga, HO, Okwu DE, Mbacbie BO. Phytochemical constituents of some Nigerian medicinal plants. *Afr. J Biotechnol.* **4**:685–688 (2005)
  31. Kavanagh F. *Analytical microbiology* Part II. (New York Academic Press, 1972) 126.